

Ultrahigh energy cosmic rays from collisional annihilation revisited

R. Dick, K.M. Hopp, and K.E. Wunderle

Abstract: We re-examine collisional annihilation of superheavy dark matter particles in dark matter density spikes in the galactic halo as a possible source of ultrahigh energy cosmic rays. We estimate the possible flux in a way that does not depend on detailed assumptions about the density profiles of dark matter clumps. The result confirms that collisional annihilation is compatible with annihilation cross sections below the unitarity bounds for superheavy dark matter if the particles can form dense cores in dark matter substructure, and it provides estimates for core sizes and densities. The ensuing clumpy source distribution in the galactic halo will be tested within a few years of operation of the Pierre Auger observatory.

Key words: Dark Matter, Ultrahigh Energy Cosmic Rays.

PACS Nos.: 98.70.Sa, 98.70.-f, 95.35.+d, 14.80.-j

1. Introduction

The source of the observed ultrahigh energy cosmic rays (UHECRs) with energies beyond the Greisen-Zatsepin-Kuzmin cutoff, $E > E_{GZK} \simeq 4 \times 10^{19}$ eV, remains a puzzle for several reasons:

1. Scattering off cosmic microwave background photons limits the penetration depths of charged particles at these energies to distances < 100 Mpc [1–3];
2. the distribution of arrival directions of UHECRs does not seem to favor any known astrophysical sources within the GZK cutoff length;
3. it seems extremely difficult to devise sufficiently efficient astrophysical acceleration mechanisms which could accelerate particles to energies $E > E_{GZK}$;

see e.g. [4–6] for recent reviews. For the second point, we note that Dolag et al. [7] find that typical deflection angles due to magnetic fields should remain below the angular resolution of the cosmic ray observatories for UHECRs originating within 100 Mpc, but the pattern of arrival directions does not seem to point to any known astrophysical accelerators in that range (but see [8] for a different opinion on the possible size of deflection angles in bottom-up models). It has been pointed out that there may be a correlation with BL Lac objects at cosmological distances [9]. However, generation in BL Lacs would require both an extremely powerful acceleration mechanism and conversion into a neutral component to avoid the GZK cutoff. This difficulty is avoided in “local” bottom-up scenarios like e.g. UHECR generation in local gamma-ray bursts within a radius of 90 Mpc [10], but Scully and Stecker

R. Dick,¹ K.M. Hopp, and K.E. Wunderle. Department of Physics and Engineering Physics, University of Saskatchewan,
116 Science Place, Saskatoon, SK S7N 5E2, Canada

¹Corresponding author (e-mail: rainer.dick@usask.ca).

had pointed out that gamma-ray bursts will not provide the observed flux [11]. Proposals of non-local explanations for absence of a GZK cutoff include a possible violation of Lorentz invariance at high energies, see [12] for a recent discussion. A recent examination of angular correlations by Gorbunov and Troitsky confirmed the absence of correlations with nearby visible objects [13].

The difficulties to explain extremely high energetic cosmic rays had motivated Hill already in 1983 to propose monopolonium decay as a possible source [14]. More recently the constraints 1-3 had also motivated Berezhinsky, Kachelrieß and Vilenkin [15] as well as Kuzmin and Rubakov [16] to propose decay of superheavy dark matter particles as sources of the observed UHECRs, see also [17]. The terrestrial flux should then be dominated by decays in our galactic dark matter halo. Decay of superheavy dark matter is an attractive proposal because it explains UHECRs from simple conversion of rest mass energy in our cosmic neighborhood, thus evading the GZK bound and the need for extremely powerful astrophysical sources. However, it is difficult to identify decay mechanisms of particles of mass $M_X \geq 10^{12}$ GeV which are slow enough to ensure a lifetime $\tau_X \geq 10^{10}$ years. This had motivated the authors of Refs. [18–20] to propose collisional annihilation of free superheavy dark matter particles (SHDM particles, WIMPZILLAS) as another mechanism to generate UHECRs in our cosmic neighborhood.

The lifetime problem is avoided in the annihilation scenario because the conversion of rest mass energy into UHECRs proceeds through two-particle collisions, thus providing a kinematic suppression of the reaction rate $\dot{n}_X \propto -n_X^2 \langle \sigma_A v \rangle$. The three UHECR puzzles are solved as in the decay scenarios: direct conversion of rest mass energy eliminates the need for an extremely powerful acceleration mechanism; the UHECRs originate within our galactic halo; and the distribution of arrival directions is determined by dense dark matter subclumps in the galactic halo, rather than by the locations e.g. of active galactic nuclei.

However, the collisional annihilation scenario suffers from its own specific shortcoming: the minuteness of the annihilation cross section $\sigma \sim M^{-2}$ of supermassive particles on the one hand ensures that these particles can survive to the present day (even with about equal amounts of matter and anti-matter), but on the other hand requires high-density substructure in the galactic halo to generate the observed flux. The results of [18] used two separate models for dark matter subhalo profiles and indicated a strong dependence of the flux on the shape of the dark matter clumps. This left open whether collisional annihilation could be a generic possibility or might be constrained by strong restrictions on subhalo profiles. We present a simplified estimate of the flux from collisional annihilation in SHDM clumps to answer this question. The main purpose of our simplified estimate is to clarify the dependence of the expected flux from SHDM annihilation on key parameters like core densities and core radii of dark matter subhalos in the galactic halo. We find that the hypothesis of collisional SHDM annihilation is compatible with the observed flux if the cores of dark matter subclumps of the galactic halo can reach densities of order $10^{-1} \rho_\odot$ and extensions of a few AU. This corresponds to a mass $\sim 10^7 M_\odot$ and would indicate that about 10% of the mass of the smallest (and most abundant) substructures detectable with N -body simulations would be bound in dense cores². We denote these dense cores as “WIMPZILLA stars”.

Besides the lifetime constraint for relic decay models and the cross section constraint for annihilation, top-down scenarios in general may be constrained by indications that the highest energy cosmic rays could be hadron dominated, see e.g. [23]. On the other hand, careful re-analysis showed that the observed highest energy event could have arisen from a primary photon [24].

Photon domination would be expected e.g. if superheavy neutrinos annihilate through an intermediate Z or a Higgs boson, because the ensuing π^0 shower would decay into UHE photons over mean free paths of order 10 km. The composition of showers in such a model would therefore resemble Z -

² Observational support for the existence of CDM subhalos within galactic halos comes from gravitational lensing [21, 22].

bursts, which assume annihilation of ultrahigh energy neutrinos (instead of supermassive neutrinos) through scattering with background neutrinos [25–27]. However, the different origin of Z -bursts implies a markedly different anisotropy signature: Z -bursts at relic neutrinos in our GZK volume would produce an isotropic pattern of arrival directions without connection to the galactic halo.

The first crucial experimental test for our top-down model of SHDM annihilation in WIMPZILLA stars will be the observed anisotropy signature after a few years of operation of the Pierre Auger observatory [28]:

If our annihilation model is correct, a pointlike source distribution should emerge with increasing source density towards the galactic center [20]. In particular, the majority of observed UHE events after about four years should appear in doublets or higher multiplets.

Alternatively, if the decay model is correct, the anisotropy signature after about five years should trace the galactic dark matter halo with uniformly increasing intensity towards the galactic center.

In Sec. 2 we discuss the generation of weakly coupled particles during inflation and explain why stable particles with masses $m \geq 10^{12}$ GeV are still around to create the observed UHE cosmic rays through annihilation. In Secs. 3 and 4 we simplify the estimates from [18] for the UHE cosmic ray flux which might be expected from SHDM annihilation in dark matter substructures within our galactic halo. The method developed in Secs. 3 and 4 yields improved estimates of typical core sizes and densities of these substructures. Sec. 3 reviews the basic formalism for the calculation of cosmic ray fluxes from dark matter annihilation, and in Sec. 4 we present our simplified estimate of the flux from SHDM annihilation. We set $\hbar = c = 1$ except in equations which are used for numerical evaluations. The differential flux per energy interval is $j(E) = dI/dE$, and the flux per unit of solid angle is $J(E) = d^2I/dEd\Omega$.

2. Superheavy dark matter from non-adiabatic expansion

One of the most interesting sources for superheavy dark matter particles is gravitational production during non-adiabatic expansion. This mechanism is usually considered in terms of the Bogolubov transformation between in and out vacua [29, 30], and was also extensively discussed as a source of superheavy dark matter particles [31–38]. Here we would like to point out how this mechanism for particle production can be understood in terms of the evolution equation for weakly coupled helicity states in an expanding universe. In the present paper we will discuss this only for weakly coupled helicity states which are not ultra-relativistic ($\lambda \not\ll m^{-1}$), although we are sure that similar conclusions can be drawn for the ultra-relativistic case, too.

The direct coupling between non-adiabaticity and changing particle numbers during gravitational expansion follows directly from the Einstein equation and the cosmological principle, since it is well known that the evolution equations for the Robertson–Walker metric

$$\frac{\dot{a}^2 + k}{a^2} = \frac{\kappa}{3}\rho \quad (1)$$

and

$$2\frac{\ddot{a}}{a} + \frac{\dot{a}^2 + k}{a^2} = -\kappa p$$

imply energy balance with only mechanical energy:

$$d(\rho a^3) = -p da^3, \quad (2)$$

and therefore the first law of thermodynamics splits into Eq. (2) and a separate balance equation for entropy and particle densities:

$$Td(sa^3) = -\sum_i \mu_i d(\nu_i a^3). \quad (3)$$

At the same time Eq. (1) usually implies a decrease or at best constancy of the comoving energy density $a^3(t)\varrho(t)$ during expansion. Consider e.g. the standard linear dispersion relation for radiation or dust

$$p = \left(\frac{\ell}{3} - 1\right) \varrho, \quad (4)$$

where the parameter is $\ell = 4$ for radiation and $\ell = 3$ for dust. After inflation, when k can be neglected and radiation initially dominates, the scale factor and dominant energy density evolve according to $a(t) \propto t^{1/2}$ and $\varrho(t) \propto t^{-2}$, leading to a decrease $a^3(t)\varrho(t) \propto t^{-1/2}$ of the comoving energy density. However, during inflation we have $p \simeq -\varrho$ leading to constant energy density ϱ and an exponential increase $a^3(t)\varrho(t) \propto \exp(3Ht)$ of the comoving energy density. From this argument we expect an energy increase at least in all those states which cannot readily redistribute their energy through sufficiently strong interactions, and inspection of the corresponding evolution equations for weakly coupled helicity states confirms this:

Towards the end of inflation the curvature parameter k is negligible on subhorizon scales, and weakly coupled helicity states have to satisfy

$$\ddot{\phi}(\mathbf{k}, t) + 3H\dot{\phi}(\mathbf{k}, t) + (m^2 + \mathbf{k}^2 \exp(-2Ht)) \phi(\mathbf{k}, t) = 0. \quad (5)$$

Here $\phi(\mathbf{k}, t)$ is defined through Fourier expansion with respect to the dimensionless comoving coordinates $\mathbf{x}$

$$\phi(\mathbf{x}, t) = \frac{1}{\sqrt{2\pi^3}} \int d^3\mathbf{k} \phi(\mathbf{k}, t) \exp(i\mathbf{k} \cdot \mathbf{x}).$$

The factor $a_1 = a(t_1) \exp(-Ht_1)$ is related to the scale factor $a(t_1)$ at the beginning of inflation. Towards the end of inflation the exponential factor in Eq. (5) is exponentially small, $\exp(-2Ht) \approx \exp(-140)$, and we find the same approximate evolution equation for all modes which are not ultra-relativistic ($|\mathbf{k}| \gg m$) at the end of inflation:

$$\ddot{\phi}(\mathbf{k}, t) + 3H\dot{\phi}(\mathbf{k}, t) + m^2\phi(\mathbf{k}, t) \simeq 0. \quad (6)$$

H is constant during inflation, and therefore the time evolution of the comoving energy density in the weakly coupled helicity states is given by

$$\begin{aligned} a^3(t)\varrho_\phi(\mathbf{k}, t) &\simeq \frac{1}{2}a^3(t) \left(\dot{\phi}(\mathbf{k}, t)\dot{\phi}(-\mathbf{k}, t) + m^2\phi(\mathbf{k}, t)\phi(-\mathbf{k}, t) \right) \\ &\simeq A_+ \exp\left(t\sqrt{9H^2 - 4m^2}\right) + A_- \exp\left(-t\sqrt{9H^2 - 4m^2}\right) + B. \end{aligned}$$

This implies a growing mode in the comoving energy density of weakly coupled states with $m < 1.5H \simeq 10^{14}$ GeV. What is special about the superheavy particles is that their comoving energy density is conserved after inflation, because the behavior of massive ($m > t^{-1}$) weakly coupled helicity states in the subsequent radiation and dust dominated backgrounds preserves their energy:

The asymptotic solution for weakly coupled massive states with $m > t^{-1}$ in such a background yields

$$\phi(\mathbf{k}, t) \propto t^{-3/\ell} \cos(mt + \varphi),$$

$$\varrho_\phi(\mathbf{k}, t) \propto t^{-6/\ell} \propto a^{-3},$$

and this implies in particular that the comoving density of massive particles freezes out at the end of inflation ($t \simeq 10^{-36}$ s) if

$$m > t^{-1} \simeq 10^{12} \text{ GeV}.$$

From these considerations follows a mass window for inflationary production of superheavy relic particles

$$10^{12} \text{ GeV} < m < 10^{14} \text{ GeV}.$$

It is certainly reassuring for the top-down models that a typical time scale for the duration of inflation yields the UHECR mass scale for the preserved particles.

3. Basic formalism

Following [18], we consider annihilation of two superheavy dark matter particles of mass $M_X \geq 10^{12} \text{ GeV}$ primarily into two jets of energy M_X .

The cosmic ray flux per energy interval originating from annihilation of particles and anti-particles of total density $n_X(\mathbf{r})$ is

$$j(E, \mathbf{r}_\odot) = \frac{dI(E, \mathbf{r}_\odot)}{dE} = \frac{d\mathcal{N}(E, M_X)}{dE} \int d^3\mathbf{r} \frac{n_X^2(\mathbf{r}) \langle \sigma_A v \rangle}{8\pi |\mathbf{r}_\odot - \mathbf{r}|^2} \quad (7)$$

if the particles are *not* Majorana particles, or $4 \times j(E, \mathbf{r}_\odot)$ otherwise. $d\mathcal{N}(E, M_X)$ is the number of particles in the energy interval $[E, E + dE]$ emerging from a jet of energy M_X and two primary jets are assumed. Neglecting events with more than two jets, $d\mathcal{N}(E, M_X)/dE$ is related to fragmentation functions via

$$\frac{d\mathcal{N}(E, M_X)}{dE} = \sum_i \frac{1}{2\sigma_A} \frac{d\sigma^{(i)}}{dE} = \frac{1}{2M_X} \sum_i F^{(i)}(x, 4M_X^2),$$

where $x = E/M_X$ and $F^{(i)}(x, 4M_X^2)$ is the differential number of particles of species i generated in the prescribed x -range in an annihilation event with $s = 4M_X^2$.

As shown in [18], UHECRs from annihilation of superheavy dark matter particles provide already a good fit to the shape of the observed spectrum if the old MLLA fragmentation function [39] is used for an estimate on $d\mathcal{N}(E, M_X)/dE$. More refined approximations have been employed in the discussion of the decay scenario [17, 40–43], and these also further improve the corresponding fit for the annihilation scenario, since for the spectral shape the transition between the decay and annihilation scenarios only corresponds to a scaling of jet energies

$$\left. \frac{d\mathcal{N}(E, M_X/2)}{dE} \right|_{\text{decay}} \rightarrow \left. \frac{d\mathcal{N}(E, M_X)}{dE} \right|_{\text{annihilation}} \quad (8)$$

in the fragmentation function.

4. A simplified estimate of the flux from dark matter clumps in the galactic halo

Numerical simulations indicate that the dark matter halos of galaxies have a lot of substructure in terms of spatial density fluctuations. In [18] it was found that the flux from local overdensities, or dark matter clumps, exceeds the flux from the smooth halo component by far. Here we will reconsider these calculations with a view on the density n_X and size r_{core} of the central core regions of dark matter clumps. n_X is of crucial importance for the UHECR flux from annihilation in these clumps.

We also want to simplify the flux estimate with regard to assumptions on the distribution of sub-clumps. In [18] a clump distribution

$$n_{cl}(d, M_{cl}) \propto \left(\frac{M_{cl}}{M_H} \right)^{-\alpha} \left[1 + \left(\frac{d}{d_0} \right)^2 \right]^{-3/2} \quad (9)$$

was assumed, where $n_{cl}(d, M_{cl})$ is the volume density of clumps with mass M_{cl} at distance d from the galactic center, d_0 is a scale radius for the subclump distribution in the galaxy, and M_H is the mass of the galactic halo [44]. Some of the recent work on structure formation has questioned the assumption of increasing clump density towards the galactic center because tidal stripping would more severely affect large clumps close to the galactic center. However, the distribution of the 150 known globular clusters in the galaxy suggests an increase of visible substructure towards the galactic center [45], and it seems reasonable to assume a similar distribution for dark matter substructure.

In the present paper, we want to avoid any detailed assumptions like (9) about the profile of the substructure distribution in the galactic halo, and we also want to avoid any assumptions about the density profiles of substructure, except for the reasonable assumption that the substructures will have dense cores with an average mass density $n_X M_X$.

The annihilation flux from an approximately compact source of volume V at distance $d(\gg V^{1/3})$ is

$$j(E, \mathbf{r}_\odot) = \int_V d^3\mathbf{r} \frac{1}{8\pi|\mathbf{r}_\odot - \mathbf{r}|^2} \frac{d\mathcal{N}(E, M_X)}{dE} n_X(\mathbf{r})^2 \langle \sigma_A v \rangle \simeq \frac{V}{8\pi d^2} \frac{d\mathcal{N}(E, M_X)}{dE} n_X^2 \langle \sigma_A v \rangle. \quad (10)$$

UHECR experiments often plot the approximately flat rescaled flux

$$E^3 j(E, \mathbf{r}_\odot) \simeq \frac{E^3 V}{8\pi d^2 M_X c^2} \frac{d\mathcal{N}(x, M_X)}{dx} n_X^2 \langle \sigma_A v \rangle,$$

where $x = E/M_X c^2$ is energy in units of primary jet energy. A typical value for fragmentation functions at $x = 0.1$ is $d\mathcal{N}(x, E_{\text{jet}})/dx \simeq 30$.

In the following we parametrize the annihilation cross section in terms of the s-wave unitarity bound for $M c^2 = 10^{12}$ GeV and $v = 100$ km/s:

$$\langle \sigma_A v \rangle = \xi \times \frac{4\pi\hbar^2}{M^2 v} = \xi \times 4.40 \times 10^{-43} \text{ m}^3/\text{s}, \quad (11)$$

where the unitarity bound is $\xi \leq 1$.

For a fiducial average distance we use $d = 10$ kpc, whereas for the fiducial volume and density of subclump cores we should have

$$n_X M_X \bar{V}_{core} \simeq 0.1 \bar{M}_{cl} = 0.1 f_{cl} M_{halo} / N_{cl} \simeq 10^{-5} M_{halo}.$$

Here f_{cl} is the mass fraction in the N_{cl} dark matter subclumps. Numerical N -body simulations and gravitational lensing indicate that f_{cl} should be of the order of a few percent. E.g. a core density

$$n_X M_X = \eta \varrho_\odot = \eta \times 1.41 \times 10^3 \text{ kg/m}^3, \\ n_X = \eta \times \frac{\varrho_\odot}{10^{12} \text{ GeV}/c^2} = \eta \times 7.89 \times 10^{17} \text{ m}^{-3},$$

yields typical core volumes and radii

$$\bar{V}_{core} \simeq \eta^{-1} \times 2 \times 10^7 V_\odot, \\ \bar{r}_{core} \simeq \eta^{-1/3} \times 1.9 \times 10^{11} \text{ m}.$$

With these fiducial average values, the flux at $E = 10^{11}$ GeV from N_{cl} dense cores of clumps can be estimated as

$$E^3 j(E) \Big|_{E=10^{11} \text{ GeV}} \simeq N_{cl} \bar{V}_{core} \frac{E^3}{8\pi d^2 M_X c^2} \frac{d\mathcal{N}(x, M_X)}{dx} n_X^2 \langle \sigma_A v \rangle \quad (12) \\ \simeq 0.1 f_{cl} M_{halo} \frac{E^3}{8\pi d^2 M_X^2 c^2} \frac{d\mathcal{N}(x, M_X)}{dx} n_X \langle \sigma_A v \rangle$$

$$\begin{aligned}
&\simeq 5.83 \times 10^{25} \text{ eV}^2 \text{ m}^{-2} \text{ s}^{-1} \times \frac{f_{cl}}{0.06} \times \frac{M_{halo}}{2 \times 10^{12} M_{\odot}} \times \left(\frac{d}{10 \text{ kpc}} \right)^{-2} \times \frac{\eta \xi}{10^{-3}}, \\
E^3 J(E) \Big|_{E=10^{11} \text{ GeV}} &\simeq 4.64 \times 10^{24} \text{ eV}^2 \text{ m}^{-2} \text{ s}^{-1} \text{ sr}^{-1} \\
&\times \frac{f_{cl}}{0.06} \times \frac{M_{halo}}{2 \times 10^{12} M_{\odot}} \times \left(\frac{d}{10 \text{ kpc}} \right)^{-2} \times \frac{\eta \xi}{10^{-3}}.
\end{aligned} \tag{13}$$

This is compatible with the UHECR flux observed by AGASA³ [46],

$$E^3 J(E) \Big|_{E=10^{11} \text{ GeV}} \simeq 4.2 \times 10^{24} \text{ eV}^2 \text{ m}^{-2} \text{ s}^{-1} \text{ sr}^{-1},$$

if the product of the parameters η and ξ is of order $\eta \xi \sim 10^{-3}$. E.g. reproduction of the beautiful standard model fragmentation fit in Fig. 7 from Ref. [41] with the present approximations would require $\eta \xi \simeq 8.6 \times 10^{-4}$.

5. Conclusion

The result (12) indicates that collisional annihilation of superheavy dark matter particles remains a viable scenario for the origin of UHECRs, albeit at the expense of postulating dense cores in dark matter substructure. This difficulty compares e.g. to the lifetime problem in decay scenarios, or to the problem of defeating synchrotron radiation loss and collisional energy loss in bottom-up scenarios.

The unitarity bound (11) on the annihilation cross section is both a virtue and a curse of the collisional annihilation scenario. It is a virtue because it implies that ultrahigh energy cosmic rays from collisional annihilation must originate in dense cores of dark matter clumps in the galactic halo, amounting to a unique prediction: an unmistakable clumpy pattern in arrival directions should emerge after four to five years of operation of the Pierre Auger observatory, appearing as pointlike sources within our halo. The connection to the halo should become apparent through an increase of source density towards the galactic center, and it will eliminate extragalactic scenarios for UHE origin. In addition, the clustering of arrival directions distinguishes it from the SHDM decay scenarios, and the absence of correlations of the pointlike sources with supernovae will rule out a bottom-up origin. A further advantage of the unitarity bound is that it ensures that stable superheavy dark matter particles will still be around due to the extremely slow rates of any reactions involving these particles and whatever affected the low mass modes to generate the baryon asymmetry after inflation very likely could not affect the superheavy modes. Yet at the same time the unitarity bound seems to require high core densities, and this may collisional annihilation look less appealing in the current stage, where it is not explicitly ruled in or ruled out. However, the required core densities for dark matter subclumps are not unreasonable and the possibility of UHECRs from collisional SHDM annihilation can not easily be dismissed.

From our point of view, primary composition is a secondary criterion for separation of different scenarios for UHECR origin. Inferring primary composition from observations requires particle physics modelling with extrapolations to very high energies whereas the interpretation of anisotropy will involve much less theoretical uncertainty.

Either way, the anisotropy signature observed by the Pierre Auger observatory will decide the fate of the annihilation scenario and we are cautiously optimistic that the collisional SHDM annihilation mechanism in WIMPZILLA stars will withstand the tests of the near future.

³The HiRes collaboration uses a different binning of the AGASA data, but still finds a similar AGASA flux at and above $E = 10^{11} \text{ GeV}$ without a cutoff, see e.g. [47]. However, the HiRes collaboration reports a cutoff in the spectrum based on its own data. This issue will be resolved by the Pierre Auger Observatory due to the much improved statistics expected from that experiment, and due to the combination of Cherenkov detectors and air fluorescence telescopes.

Acknowledgement

This research was supported in part by the Natural Sciences and Engineering Research Council of Canada. We are grateful to Floyd Stecker for helpful comments on an early draft of this paper.

References

1. K. Greisen, Phys. Rev. Lett. **16**, 748 (1966); G.T. Zatsepin, V.A. Kuzmin, JETP Lett. **4**, 78 (1966).
2. F.W. Stecker, Phys. Rev. Lett. **21**, 1016 (1968).
3. T. Stanev, R. Engel, A. Mücke, R.J. Protheroe, J.P. Rachen, Phys. Rev. **D62**, 093005 (2000).
4. D.F. Torres, L.A. Anchordoqui, Rep. Prog. Phys. **67**, 1663 (2004).
5. F.W. Stecker, J. Phys. **G29**, R47 (2003).
6. T.K. Gaisser, Nucl. Phys. B (Proc. Suppl.) **117**, 318 (2003).
7. K. Dolag, D. Grasso, V. Springel, I. Tkachev, JETP Lett. **79**, 583 (2004); JCAP **0501**, 009 (2005).
8. G. Sigl, F. Miniati, T. Ensslin, Phys. Rev. **D70**, 043007 (2004), Nucl. Phys. B (Proc. Suppl.) **136**, 224 (2004).
9. P.G. Tinyakov, I.I. Tkachev, JETP Lett. **74**, 445 (2001).
10. E. Waxman, Phys. Rev. Lett. **75**, 386 (1995), *Gamma-ray bursts: Potential sources of ultra high energy cosmic rays*, astro-ph/0412554.
11. S.T. Scully, F.W. Stecker, Astropart. Phys. **16**, 271 (2002).
12. F.W. Stecker, S.T. Scully, Astropart. Phys. **23**, 203 (2005).
13. D.S. Gorbunov, S.V. Troitsky, Astropart. Phys. **23**, 175 (2005).
14. C.T. Hill, Nucl. Phys. **B224**, 469 (1983).
15. V. Berezhinsky, M. Kachelrieß, A. Vilenkin, Phys. Rev. Lett. **79**, 4302 (1997).
16. V.A. Kuzmin, V.A. Rubakov, in *Beyond the Desert — Accelerator and Non-Accelerator Approaches*, edited by H.V. Klapdor-Kleingrothaus and H. Päs, IoP Publishing, Bristol 1998, pp. 802–807; Phys. Atom. Nucl. **61**, 1028 (1998).
17. M. Birkel, S. Sarkar, Astropart. Phys. **9**, 297 (1998).
18. P. Blasi, R. Dick, E.W. Kolb, Astropart. Phys. **18**, 57 (2002).
19. P. Blasi, R. Dick, E.W. Kolb, Nucl. Phys. B (Proc. Suppl.) **110**, 494 (2002).
20. R. Dick, P. Blasi, E.W. Kolb, Nucl. Phys. B (Proc. Suppl.) **124**, 201 (2003).
21. M. Chiba, Astrophys. J. **565**, 17 (2002).
22. N. Dalal, C.S. Kochanek, Astrophys. J. **572**, 25 (2002).
23. A.A. Watson, *The mass composition of cosmic rays above 10^{17} eV*, astro-ph/0410514.
24. P. Homola et al., Acta Phys. Pol. **B35**, 1893 (2004); M. Risse et al., *On the primary particle type of the most energetic Fly's Eye experiment*, astro-ph/0410739.
25. T. Weiler, Astropart. Phys. **11**, 303 (1999); D. Fargion, B. Mele, A. Salis, Astrophys. J. **517**, 725 (1999).
26. Z. Fodor, S. Katz, A. Ringwald, JHEP **0206**, 046 (2002).
27. G. Gelmini, G. Varieschi, T. Weiler, Phys. Rev. **D70**, 113005 (2004).
28. K.-H. Kampert (for the Pierre Auger Collaboration), *The Pierre Auger Observatory - Status and Prospects*, astro-ph/0501074.
29. L. Parker, Phys. Rev. **183**, 1057 (1969).
30. N.D. Birrell, P.C.W. Davies, *Quantum Fields in Curved Space*, Cambridge University Press, Cambridge 1982.
31. J.D. Barrow, E.J. Copeland, E.W. Kolb, A.R. Liddle, Phys. Rev. **D43**, 977 (1991).
32. A. Masiero, A. Riotto, Phys. Lett. **B289**, 73 (1992).
33. E.W. Kolb, A. Riotto, I.I. Tkachev, Phys. Lett. **B423**, 348 (1998).
34. D.J.H. Chung, E.W. Kolb, A. Riotto, Phys. Rev. **D59**, 023501 (1999), Phys. Rev. **D60**, 063504 (1999).
35. V.A. Kuzmin, I.I. Tkachev, Phys. Rev. **D59**, 123006 (1999), Phys. Rep. **320**, 199 (1999).
36. G.F. Giudice, M. Peloso, A. Riotto, I.I. Tkachev, JHEP **9908**, 014 (1999).
37. D.J.H. Chung, E.W. Kolb, A. Riotto, I.I. Tkachev, Phys. Rev. **D62**, 043508 (2000).
38. D.J.H. Chung, P. Crotty, E.W. Kolb, A. Riotto, Phys. Rev. **D64**, 043503 (2001).
39. Ya.I. Azimov, Yu.L. Dokshitser, V.A. Khoze, S.I. Troyan, Phys. Lett. **B165**, 147 (1985), Z. Phys. **C27**, 65 (1985), Z. Phys. **C31**, 213 (1986).

40. Z. Fodor, S.D. Katz, Phys. Rev. Lett. **86**, 3224 (2001).
41. S. Sarkar, R. Toldrà, Nucl. Phys. **B621**, 495 (2002).
42. C. Barbot, M. Drees, Astropart. Phys. **20**, 5 (2003).
43. R. Aloisio, V. Berezhinsky, M. Kachelrieß, Phys. Rev. **D69**, 094023 (2004).
44. P. Blasi, R. Sheth, Phys. Lett. **B486**, 233 (2000).
45. W.E. Harris, Astron. J. **112**, 1487 (1996).
46. M. Takeda et al., Astropart. Phys. **19**, 447 (2003).
47. R.U. Abbasi et al., Phys. Rev. Lett. **92**, 151101 (2004).